arXiv:1207.2436v1 [math.CA] 7 Jul 2012

NEW ESTIMATES ON INTEGRAL INEQUALITIES AND THEIR APPLICATIONS

M.EMIN ÖZDEMIR[♦], MUSTAFA GÜRBÜZ[♣], AND MEVLÜT TUNÇ[♣]

Abstract. In this paper, we obtain some inequalities by using a kernel and an inequality which is a result of Young inequality. Besides we give some applications to special means.

1. Introduction

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function on the interval I of real numbers and $a, b \in I$ with $a < b$. The inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)\,dx \leq \frac{f(a)+f(b)}{2} \tag{1.1}$$

is well known in the literature as Hermite-Hadamard's inequality for convex functions [1].

In [2], Dragomir and Agarwal proved one lemma and some Hadamard's type inequalities for convex functions as following:

Lemma 1. *Let $f : I^\circ \subseteq \mathbb{R} \to \mathbb{R}$ be a differantiable mapping on I°, $a, b \in I^\circ$ with $a < b$. If $f' \in L[a, b]$, then the following equality holds:*

$$\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx = \frac{b-a}{2}\int_0^1 (1-2t)\, f'(ta + (1-t)\,b)\,dt.$$

Theorem 1. *Let $f : I^\circ \subseteq \mathbb{R} \to \mathbb{R}$ be a differantiable mapping on I°, $a, b \in I^\circ$ with $a < b$. If $|f'|$ is convex on $[a, b]$, then the following inequality holds:*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx\right| \leq \frac{(b-a)\left(|f'(a)| + |f'(b)|\right)}{8}.$$

Theorem 2. *Let $f : I^\circ \subseteq \mathbb{R} \to \mathbb{R}$ be a differantiable mapping on I°, $a, b \in I^\circ$ with $a < b$, and let $p > 1$. If the new mapping $|, f'|^{p/(p-1)}$ is convex on $[a, b]$, then the following inequality holds:*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)\,dx\right| \leq \frac{(b-a)}{2(p+1)^{1/p}}\left[\frac{\left(|f'(a)|^{p/(p-1)} + |f'(b)|^{p/(p-1)}\right)}{2}\right]^{(p-1)/p}.$$

We recall the well-known Young's inequality which can be stated as follows.

Key words and phrases. Young inequality, hölder inequality, power-mean inequality.

Theorem 3. *(**Young's inequality**, see* [3], *p. 117) If $a, b > 0$ and $p, q > 1$ satisfy $\frac{1}{p} + \frac{1}{q} = 1$, then*

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}. \tag{1.2}$$

Equality holds if and only if $a^p = b^q$.

Remark 1. [4] *If we take $a = t^{\frac{1-p}{p^2}}$ and $b = t^{\frac{1}{pq}}$ in (1.2), we have*

$$1 \leq \frac{1}{p} t^{\frac{1}{p}-1} + \left(1 - \frac{1}{p}\right) t^{\frac{1}{p}} \tag{1.3}$$

for all $t \in [0, 1]$.

In the following sections our main results are given.

2. New Results

Theorem 4. *Let $f : I^\circ \subseteq \mathbb{R} \to \mathbb{R}$ be a differantiable convex mapping on I°, $a, b \in I^\circ$ with $a < b$. If $f, f' \in L[a, b]$, for $p, q > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, the following inequality holds:*

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)\, dx \right| \leq \frac{p}{(p+1)^{1+\frac{1}{p}}} \left(\int_a^b |f'(x)|^q\, dt \right)^{\frac{1}{q}}$$

Proof. By Lemma 1 in [2] we have

$$\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)\, dx = \frac{b-a}{2} \int_0^1 (1-2t) f'(ta + (1-t) b)\, dt. \tag{2.1}$$

As we choose f is convex on I° by using Hadamard's inequality, we can see that both sides are positive of Lemma 2.1

On the other hand by using Young's inequality we have ($t \in [0, 1]$, $p > 1$)

$$1 \leq \frac{1}{p} t^{\frac{1}{p}-1} + \left(1 - \frac{1}{p}\right) t^{\frac{1}{p}}$$

which is proved in [4]. If we integrate both sides of above inequality respect to t over $[0, 1]$ we get

$$1 \leq \int_0^1 \left(\frac{1}{p} t^{\frac{1}{p}-1} + \left(1 - \frac{1}{p}\right) t^{\frac{1}{p}} \right) dt \tag{2.2}$$

By multipliying both sides of (2.1) and (2.2) we have

$$\begin{aligned} \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)\, dx \quad &\leq \quad \frac{b-a}{2} \int_0^1 \left(\frac{1}{p} t^{\frac{1}{p}-1} + \left(1 - \frac{1}{p}\right) t^{\frac{1}{p}} \right) dt \\ &\quad \times \int_0^1 (1-2t) f'(ta + (1-t) b)\, dt \end{aligned} \tag{2.3}$$

To use Hölder's inequality we apply properties of ablosute value as

$$\begin{aligned}\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| &\le \frac{b-a}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)dt\\ &\quad\times\int_0^1\left|(1-2t)\,f'(ta+(1-t)\,b)\right|dt\end{aligned}$$

By using Hölder's inequality we have

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\\ \le\;& \frac{b-a}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)dt\\ &\times\left(\int_0^1|1-2t|^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1|f'(ta+(1-t)\,b)|^q\,dt\right)^{\frac{1}{q}}\\ =\;& \frac{b-a}{2}\left(\frac{2p}{p+1}\right)\left(\int_0^{\frac{1}{2}}(1-2t)^p\,dt+\int_{\frac{1}{2}}^1(2t-1)^p\,dt\right)^{\frac{1}{p}}\\ &\times\left(\int_0^1|f'(ta+(1-t)\,b)|^q\,dt\right)^{\frac{1}{q}}\\ =\;& \frac{b-a}{2}\left(\frac{2p}{p+1}\right)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\int_0^1|f'(ta+(1-t)\,b)|^q\,dt\right)^{\frac{1}{q}}.\end{aligned}$$

By simple calculation we get the desired result. □

Corollary 1. *If we choose $p=q=2$ in Theorem 4, we have*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\le\frac{2}{3^{\frac{3}{2}}}\left(\int_a^b|f'(x)|^2\,dt\right)^{\frac{1}{2}}.$$

Theorem 5. *Let $f: I^\circ\subseteq\mathbb{R}\to\mathbb{R}$ be a differantiable convex mapping on I°, $a,b\in I^\circ$ with $a<b$. If $f,f'\in L[a,b]$, for $p,q>1$, $\frac{1}{p}+\frac{1}{q}=1$, the following inequality holds:*

$$\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\le\frac{p(p-1)}{(p+1)(2p+1)}\left[f(b)-f(a)\right]$$

Proof. The same steps are followed as in Theorem 4 until (2.3). Then we know that the function

$$f(t)=1-2t$$

is nonincreasing on $[0,1]$, and since f is convex, f'' is positive on I°. So f' is nondecreasing. By using these phrases we can use Chebyshev inequality as:

$$\begin{aligned}
& \frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx \\
\leq\ & \frac{b-a}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)dt \\
& \times\int_0^1 (1-2t)\,f'(ta+(1-t)b)\,dt \\
\leq\ & \frac{b-a}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)dt \\
& \times\int_0^1 (1-2t)\,dt\int_0^1 f'(ta+(1-t)b)\,dt \\
=\ & \frac{f(b)-f(a)}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)dt \\
& \times\int_0^1 (1-2t)\,dt
\end{aligned}$$

Since both of the functions $\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)$ and $(1-2t)$ are nonincreasing, we can use Chebyshev inequality again as:

$$\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\leq\frac{f(b)-f(a)}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)(1-2t)\,dt$$

By simple calculation, the proof is completed. ☐

Corollary 2. *If we choose $p=1,1$ in Theorem 5, we have*

$$\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\leq\frac{11}{483}\left[f(b)-f(a)\right].$$

Theorem 6. *Let $f: I^\circ\subseteq\mathbb{R}\to\mathbb{R}$ be a differantiable convex mapping on I°, $a,b\in I^\circ$ with $a<b$. If $f,f'\in L[a,b]$, for $p,q>1$, $\frac{1}{p}+\frac{1}{q}=1$, the following inequality holds:*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\leq\frac{2^{\frac{1}{q}}p}{(p+1)(b-a)}\left(\int_a^b\left|x-\frac{a+b}{2}\right|\left|f'(x)\right|^q dx\right)^{\frac{1}{q}}.$$

Proof. The same steps are followed as in Theorem 4 until (2.3). Then by using convexity of f and properties of absolute value we have

$$\begin{aligned}
\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \leq\ & \frac{b-a}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)dt \\
& \times\int_0^1\left|(1-2t)\,f'(ta+(1-t)b)\right|dt.
\end{aligned}$$

By using Power-mean inequality we have

$$\begin{aligned}(2.4)\qquad & \left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right| \\ \leq\ & \frac{b-a}{2}\int_0^1\left(\frac{1}{p}t^{\frac{1}{p}-1}+\left(1-\frac{1}{p}\right)t^{\frac{1}{p}}\right)dt \\ & \times\left(\int_0^1|1-2t|\,dt\right)^{\frac{1}{p}}\left(\int_0^1|(1-2t)|\,|f'(ta+(1-t)b)|^q\,dt\right)^{\frac{1}{q}} \\ =\ & \frac{2^{\frac{1}{q}}p}{p+1}\left(\int_0^1|(1-2t)|\,|f'(ta+(1-t)b)|^q\,dt\right)^{\frac{1}{q}}\end{aligned}$$

And using the change of the variable $x=ta+(1-t)b$, $t\in[0,1]$, inequality (2.4) can be writen as

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\leq\frac{2^{\frac{1}{q}}p}{(p+1)(b-a)}\left(\int_a^b\left|x-\frac{a+b}{2}\right||f'(x)|^q\,dt\right)^{\frac{1}{q}}.$$

Then the proof is completed. □

Corollary 3. *If we choose $p=q=2$ in Theorem 6, we have*

$$\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)\,dx\right|\leq\frac{2^{\frac{3}{2}}}{3}\left(\int_0^1\left|x-\frac{a+b}{2}\right||f'(x)|^q\,dx\right)^{\frac{1}{2}}$$

3. APPLICATIONS TO SPECIAL MEANS

We now consider the applications of our Theorems to the following special means
The arithmetic mean: $A=A(a,b):=\frac{a+b}{2},\ \ a,b\geq 0,$
The geometric mean: $G=G(a,b):=\sqrt{ab},\ \ a,b\geq 0,$
The harmonic mean: $H=H(a,b):=\frac{2ab}{a+b},\ \ a,b\geq 0,$
The logarithmic mean: $L=L(a,b):=\begin{cases} a & \text{if}\quad a=b \\ \frac{b-a}{\ln b-\ln a} & \text{if}\quad a\neq b\end{cases},\ \ a,b\geq 0,$
The Identric mean: $I=I(a,b):=\begin{cases} a & \text{if}\quad a=b \\ \frac{1}{e}\left(\frac{b^b}{a^a}\right)^{\frac{1}{b-a}} & \text{if}\quad a\neq b\end{cases},\ \ a,b\geq 0,$
The p-logarithmic mean:$L_p=L_p(a,b):=\begin{cases}\left[\frac{b^{p+1}-a^{p+1}}{(p+1)(b-a)}\right]^{1/p} & \text{if}\quad a\neq b \\ a & \text{if}\quad a=b\end{cases},$
$p\in\mathbb{R}\backslash\{-1,0\};\ \ a,b>0.$

The following inequality is well known in the literature:

$$H\leq G\leq L\leq I\leq A\leq K$$

It is also known that L_p is monotonically increasing over $p\in\mathbb{R}$, denoting $L_1=A$, $L_0=I$ and $L_{-1}=L$.

The following propositions holds:

Proposition 1. *Let a,b* $\in \mathbb{R}^+$, $a < b$ *and* $n \in \mathbb{N}$, $n \geq 2$. *Then, we have*

$$|A(a^n, b^n) - L_n(a,b)| \leq \frac{n.p.(b-a)^{\frac{1}{q}}}{(p+1)^{1+\frac{1}{p}}} A^{n-1}(a,b). \tag{3.1}$$

Proof. The proof is immediate from Theorem 4 applied for $f(x) = x^n$, $x \in \mathbb{R}$. □

Proposition 2. *Let a,b* $\in \mathbb{R}^+$, $a < b$ *and* $n \in \mathbb{N}$, $n \geq 2$. *Then, for all* $p > 1$, *the following inequality holds:*

$$A(a^n, b^n) - L_n(a,b) \leq \frac{p(p-1)}{(p+1)(2p+1)} [b^n - a^n]. \tag{3.2}$$

Proof. The proof is immediate from Theorem 5 applied for $f(x) = x^n$, $x \in \mathbb{R}$. □

Proposition 3. *Let a,b* $\in \mathbb{R}^+$, $a < b$ *and* $n \in \mathbb{N}$, $n \geq 2$. *Then, for all* $p > 1$, *the following inequality holds:*

$$|A(a^n, b^n) - L_n(a,b)| \leq \frac{2^{\frac{1}{q}} p}{(p+1)(b-a)} \left(\int_a^b \left| x - \frac{a+b}{2} \right| |f'(x)|^q \, dx \right)^{\frac{1}{q}}. \tag{3.3}$$

Proof. The proof is immediate from Theorem 6 applied for $f(x) = x^n$, $x \in \mathbb{R}$. □

Proposition 4. *Let a,b* $\in \mathbb{R}^+$, $a < b$. *Then, we have*

$$\left| A\left(a^{-1}, b^{-1}\right) - L^{-1}(a,b) \right| \leq \frac{p}{(p+1)^{1+\frac{1}{p}}} \left(\int_a^b \left| -x^{-2} \right|^q dt \right)^{\frac{1}{q}}.$$

Proof. The proof is obvious from Theorem 4 applied for $f(x) = 1/x$, $x \in [a,b]$. □

Proposition 5. *Let a,b* $\in \mathbb{R}^+$, $a < b$. *Then, we have*

$$A\left(a^{-1}, b^{-1}\right) - L^{-1}(a,b) \leq \frac{p(p-1)}{(p+1)(2p+1)} \left(\frac{1}{b} - \frac{1}{a} \right).$$

Proof. The proof is obvious from Theorem 5 applied for $f(x) = 1/x$, $x \in [a,b]$. □

Proposition 6. *Let a,b* $\in \mathbb{R}^+$, $a < b$. *Then, we have*

$$\left| A\left(a^{-1}, b^{-1}\right) - L^{-1}(a,b) \right| \leq \frac{2^{\frac{1}{q}} p}{(p+1)(b-a)} \left(\int_a^b \left| x - \frac{a+b}{2} \right| \left| -x^{-2} \right|^q dx \right)^{\frac{1}{q}}.$$

Proof. The proof is obvious from Theorem 6 applied for $f(x) = 1/x$, $x \in [a,b]$. □

REFERENCES

[1] S.S. Dragomir, C. E. M. Pearce, Selected Topic on Hermite- Hadamard Inequalities and Applications, URL:http://www.maths.adelaide.edu.au /Applied /staff /cpearce.html

[2] S.S. Dragomir, R.P. Agarwal, Two Inequalities for Differentiable Mappings and Applications to Special Means of Real Numbers and to Trapezoidal Formula, Appl. Math. Lett., Vol. 11, No. 5, pp. 91-95, 1998.

[3] S.S. Dragomir, R.P. Agarwal and N.S. Barnett, Inequalities for beta and gamma functions via some classical and new integral inequalities, J. Inequal. & Appl., 5 (2000), 103-165.

[4] M. Tunç, Two New Definitions on Convexity and Related Inequalities, arXiv:1205.5189v1 [math.CA], 23 May 2012.

♦ATATÜRK UNIVERSITY, K.K. EDUCATION FACULTY, DEPARTMENT OF MATHEMATICS, ERZURUM 25240, TURKEY
E-mail address: `emos@atauni.edu.tr`

♠AĞRI İBRAHIM ÇEÇEN UNIVERSITY, FACULTY OF EDUCATION, DEPARTMENT OF MATHEMATICS, AĞRI 04100, TURKEY
E-mail address: `mgurbuz@agri.edu.tr`

♣KILIS 7 ARALIK UNIVERSITY, FACULTY OF SCIENCE AND ARTS, DEPARTMENT OF MATHEMATICS, KILIS 79100, TURKEY
E-mail address: `mevluttunc@kilis.edu.tr`